



## The Deccan beyond the plume hypothesis

H. Sheth (1)

Indian Institute of Technology, Powai, Mumbai 400 076 India (hcsbeth@iitb.ac.in)

The widely accepted mantle plume model postulates that (i) the currently volcanically active Réunion Island, in the Indian Ocean, is fed by the narrow “tail” of a mantle plume that rises from the core-mantle boundary, (ii) the Deccan continental flood basalt province, of India, originated from the “head” of the same plume during its early eruptive phase near the end of the Cretaceous, and (iii) the Lakshadweep-Chagos Ridge, an important linear volcanic ridge in the Indian Ocean, is a product of the plume. It is not generally appreciated, however, that this so-called “classic” case of a plume contradicts the plume model in many ways. For example, there is little petrological evidence as yet that the Deccan source was “abnormally hot”, and the short (~1.0-0.5 m.y.) duration claimed by some for the eruption of the Deccan is in conflict with recent Ar-Ar age data that suggest the total duration to have been at least ~8 m.y. The Deccan CFB was associated with the breakup of the Seychelles microcontinent from India. Geological and geophysical data from the Deccan provide no support to the plume model and arguably undermine it altogether. There is solid field evidence that there was no regional pre-volcanic domal lithospheric uplift in the Deccan, though such uplift is expected from a thermal plume. On the other hand, the Deccan region has undergone kilometre-scale post-volcanic uplift. The interplay of several intersecting continental rift zones in India is apparently responsible for the roughly circular outcrop of the Deccan. The Lakshadweep-Chagos Ridge, and the islands of Mauritius and Réunion, are located along fracture zones, and the apparent systematic age progression along the Ridge may be a result of southward crack propagation through the oceanic lithosphere. This idea avoids the problem of a 10-degree palaeolatitude discrepancy which the plume model can only solve with the ad hoc inclusion of mantle roll. Published Ar-Ar age data for the Lakshadweep-Chagos Ridge basalts have been seriously questioned. “Enriched” isotopic ratios such as higher-than-N-MORB values of  $^{87}\text{Sr}/^{86}\text{Sr}$ , observed in basalts of the Ridge and the Mascarene Islands may mark the involvement of delaminated enriched continental mantle instead of a plume. High

values of the  $^3\text{He}/^4\text{He}$  ratio also do not represent a deep mantle component or plume. The three Mascarene Islands (Mauritius, Réunion, and Rodrigues) are not related to the Deccan but reflect the recent (post-10 Ma) tectonic- magmatic development of the African Plate. I relate CFB volcanism to continental rifting, which often (but not always) evolves into full-fledged sea-floor spreading. I ascribe the rifting itself not to mantle plume heads but to large-scale plate dynamics themselves, possibly aided by long-term thermal insulation beneath a supercontinent which may have surface effects similar to those predicted for “plume incubation” models. Non-plume, plate tectonic models are capable of explaining the Deccan in all its greatness, and there is no trace of a mantle plume in this vast region.